

Title: Informed Consent System & Method Therefor

Inventor(s): Laurysen et al.

Appln. No. 09/782,128




Figure 1

Title: Informed Consent System & Method
Therefor

Inventor(s): Laurysen et al.

Appl. No. 09/782,128



FIGURE 2A

Title: Informed Consent System & Method
Therefor

Inventor(s): Laurysen et al.

Appl. No. 09/782,128



FIGURE 2B

Title: Informed Consent System & Method Therefor

Inventor(s): Laurysen et al.

Appl. No. 09/782,128




Figure 3

Title: Informed Consent System & Method Therefor
Inventor(s): Laurysen et al.
Appl. No. 09/782,128




Figure 4

Title: Informed Consent System & Method
Therefor

Inventor(s): Laurysen et al.

Appl. No. 09/782,128



Complication 13 of 17

50

Hoarseness after surgery occurs in up to 15% of (or 15 of every 100) patients and is usually caused by normal tissue swelling. However, in some patients (0.5%-4% or 5 to 40 of every 1000) hoarseness is caused by injury to one of the nerves controlling the vocal cords and may be permanent. Normal swelling is the most common cause of hoarseness after surgery to remove a cervical disk, and the hoarseness usually improves over several days without any treatment



Click here if you would like to have this complication "flagged" for further discussion with your physician or physician's representative.

54

Show that you've read the possibility of this complication

52

by entering your initials

58

and the last 4 digits of your social security number or birth date

60

then click

56

Figure 5

2009 RELEASE UNDER E.O. 14176